Mimrylo™ rusfertide for injection



Hypothetical patient.

Getting ready for your PV appointment

The following questions will take about 5 minutes to complete. Print and complete your responses and bring them to your next appointment.

1 Symptom identification

QUESTION 1

Which of the following PV symptoms have you experienced in the past few months?

Select all that apply

- ☐ Fatigue or low energy
- ☐ Itching (especially after bathing)
- ☐ Headaches, light-headedness or vision problems
- ☐ Night sweats
- ☐ Trouble concentrating or "brain fog"
- ☐ Easy bruising, nosebleeds or gum bleeding
- ☐ Bone pain
- ☐ Abdominal discomfort or fullness
- ☐ Other: _____

2 Symptom frequency

QUESTION 2

How often are you experiencing those PV symptoms?

Select one

- ☐ Every day
- ☐ More than once per week
- ☐ Less than once per week
- ☐ Less than once per month

Mimrylo™ rusfertide for injection



Hypothetical patient.

3 Impact on my daily life

QUESTION 3

How much would you say your PV symptoms impact your daily life?

1 — 2 — 3 — 4 — 5

1 = Not at all

5 = A lot

QUESTION 4

What aspects of your life does your PV and/or PV treatment have an impact on?

Select all that apply

- ☐ Energy levels
- ☐ Mood or emotional well-being
- ☐ Ability to work or do my usual activities
- ☐ Social or family life
- ☐ Sleep
- ☐ Other: _____

4 My PV treatment

QUESTION 5

Compared with your last visit, how well do you feel you are managing your PV and PV symptoms?

Select one

- ☐ Worse
- ☐ Better
- ☐ About the same
- ☐ Not sure

QUESTION 6

How well do you feel your PV is controlled with your current treatment plan?

1 — 2 — 3 — 4 — 5

1 = Poorly controlled

5 = Well controlled

Mimrylo™ rusfertide for injection



Hypothetical patient.

QUESTION 7

What, if any, concerns do you have with your current PV treatment?

Select all that apply

- ☐ Side effects
- ☐ Continued need for phlebotomies
- ☐ Symptoms not improving
- ☐ Trouble keeping up with treatment
- ☐ Logistics of getting to phlebotomy clinic
- ☐ Doesn't seem to help with my blood counts (labs)
- ☐ No current concerns
- ☐ Other: _____

QUESTION 8

How satisfied are you with your current treatment? Are you looking to try something else like MIMRYLO?

1 — 2 — 3 — 4 — 5

1 = Not satisfied and curious

5 = Very satisfied and not interested

5 My treatment goals

QUESTION 9

What are your goals for your PV treatment?

Select all that apply

- ☐ Reduction in therapeutic phlebotomies
- ☐ Reduce risk of blood clots / thromboembolic events
- ☐ Consistent hematocrit control
- ☐ Improvement in PV symptoms
- ☐ Regulate iron deficiency
- ☐ Reduction in reliance on cytoreductive therapies (eg, hydroxyurea)
- ☐ Other: _____

Mimrylo™ rusfertide for injection



Hypothetical patient.

What is MIMRYLO (rusfertide)?

MIMRYLO is a prescription medicine that is used to treat erythrocytosis (high red blood cell count) in adults with polycythemia vera.

It is not known if MIMRYLO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Before using MIMRYLO, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. MIMRYLO may harm your unborn baby.

  Females who are able to become pregnant:

  - Your healthcare provider may do a pregnancy test before you start treatment with MIMRYLO.
  - Use effective birth control (contraception) during treatment and for at least 30 days after the final dose of MIMRYLO.
  - Tell your healthcare provider right away if you become pregnant or think that you may be pregnant during treatment with MIMRYLO.
  - Stop taking MIMRYLO if you become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if MIMRYLO passes into your breast milk. Do not breastfeed during treatment with MIMRYLO and for 30 days after the final dose of MIMRYLO. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of MIMRYLO?

MIMRYLO may cause serious side effects, including:

- **New or worsening increased blood platelets (thrombocytosis)**. MIMRYLO may increase blood platelet counts. Your healthcare provider will do blood tests when you start and during treatment with MIMRYLO to check your platelet counts.
- **Injection site reactions.** MIMRYLO may cause injection-site reactions such as redness, itching, pain, swelling, bruising, hardening at the injection site, and irritation. If you get an injection site reaction, apply ice packs, steroid cream, or take medicines as needed to treat injection site pain and swelling.

Your healthcare provider may change your dose or stop treatment with MIMRYLO if you have side effects.

The most common side effects of MIMRYLO are:

- injection site reactions
- low red blood cell count (anemia)

These are not all of the possible side effects of MIMRYLO. Call your doctor for medical advice about side effects. **You may report side effects to Takeda at 1-844-662-8532 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Please see Patient Information in the MIMRYLO (rusfertide) full Prescribing Information at mimrylo.com/PI.

Takeda

TAKEDA and Takeda are registered trademarks of Takeda Pharmaceutical Company Limited.
MIMRYLO and ——— are trademarks of Takeda Pharmaceuticals U.S.A., Inc.
©2026 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 11/26 US-MIM-0122